

Lung ultrasound in acute and chronic heart failure: a clinical consensus statement of the European Association of Cardiovascular Imaging (EACVI)

Luna Gargani ^{1*}, Nicolas Girerd ², Elke Platz ³, Pierpaolo Pellicori ⁴,
Ivan Stankovic ⁵, Alberto Palazzuoli⁶, Emanuele Pivetta ^{7,8},
Marcelo Haertel Miglione ^{9,10}, Hatem Soliman-Aboumarie ^{11,12},
Eustachio Agricola ¹³, Giovanni Volpicelli ¹⁴, Susanna Price ¹⁵,
Erwan Donal ¹⁶, Bernard Cosyns ¹⁷, and Aleksandar N. Neskovic ⁵

¹Department of Surgical, Medical and Molecular Pathology and Critical Care Medicine, University of Pisa, via Paradisa 2 5614, Pisa, Italy; ²Université de Lorraine, Centre d'Investigations Cliniques Plurithématique 1433, Institut Lorrain du Cœur et des Vaisseaux Louis Mathieu, CHRU de Nancy, INSERM DCAC, F-CRIN INI-CRCT, Nancy, France; ³Cardiovascular Division, Brigham and Women's Hospital and Harvard Medical School, Boston, USA; ⁴School of Cardiovascular and Metabolic Health, University of Glasgow, Glasgow, UK; ⁵Clinical Hospital Centre Zemun, Faculty of Medicine, University of Belgrade, Serbia; ⁶Cardiovascular Diseases Unit, Cardio-Thoracic and Vascular Department, Le Scotte Hospital, University of Siena, Italy; ⁷Medicina d'Urgenza-MECAU, Presidio Molinette, A.O.U. Città della Salute e della Scienza di Torino, Italy; ⁸Department of Medical Sciences, University of Turin, Turin, Italy; ⁹EcoHaertel - Hospital Mae de Deus, Porto Alegre, Brazil; ¹⁰Federal University of Health Sciences of Porto Alegre, Porto Alegre, Brazil; ¹¹Department of Cardiothoracic Anaesthesia and Critical Care, Harefield Hospital, Royal Brompton and Harefield Hospitals, Guy's and St Thomas NHS Foundation Trust, London, UK; ¹²School of Cardiovascular Medicine and Sciences, King's College London, UK; ¹³Cardiovascular Imaging Unit, Cardio-Thoracic-Vascular Department, San Raffaele Hospital, Vita-Salute University, Milan, Italy; ¹⁴Department of Emergency Medicine, San Luigi Gonzaga University Hospital, Torino, Italy; ¹⁵Departments of Cardiology & Intensive Care, Royal Brompton & Harefield Hospitals, Guy's and St Thomas NHS Foundation Trust, National Heart and Lung Institute, Imperial College London, London, UK; ¹⁶University of Rennes, CHU Rennes, Inserm, Rennes, France; and ¹⁷Department of Cardiology, Universitair Ziekenhuis Brussel, Jette, Brussels, Belgium

Received 4 July 2023; accepted 5 July 2023; online publish-ahead-of-print 14 July 2023

Reviewers

This document was reviewed by members of the 2020–2022 EACVI Scientific Documents Committee: Magnus Bäck, Philippe B. Bertrand, Marc Dweck, Niall Keenan, and Leyla Elif Sade.

Keywords

lung ultrasound • acute heart failure • chronic heart failure • heart failure • B-lines • point-of-care ultrasound • acute dyspnoea • chest sonography

Introduction

Lung ultrasound (LUS) was introduced to intensive care units and emergency departments more than 20 years ago, primarily as a tool for the assessment of patients with acute dyspnoea.^{1,2} Since then, it has gained popularity as a quick point-of-care examination enabling clinicians to answer crucial clinical questions. Over the last decade, the cardiology community has acknowledged the potential of LUS and expanded its use further, to assist with the diagnosis and management of patients with heart failure (HF).^{3–5}

The importance of recognizing and treating pulmonary congestion is a cornerstone in the management of patients with HF.⁶ LUS is a versatile, high sensitivity point-of-care examination to detect pulmonary de-aeration due to increased extravascular lung water. It has many advantages to the extent that an integrated cardiopulmonary ultrasound

exam is likely to become the reference standard in HF care. This approach allows the aetiology of HF to be defined, through the assessment of cardiac structure and function by echocardiography, at the same time as the assessment of pulmonary congestion provided by LUS. In addition, it facilitates the exclusion of other highly prevalent conditions that may mimic/overlap with HF (e.g. pneumonia, acute lung injury/acute respiratory distress syndrome [ALI/ARDS], and pneumothorax).

LUS in HF: the findings

B-lines and pleural effusions

In a fully aerated lung, the only anatomical structure that can be visualized is the pleura, which appears as a smooth, hyperechoic horizontal line that moves synchronously with respiration. This line is called

* Corresponding author. E-mail: luna.gargani@unipi.it

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

SHOUT
SHARE IT



'pleural line' and its movement is the 'lung sliding', providing visual assessment of lung excursion during ventilation. The sonographic pattern of the aerated lung also includes parallel, hyperechoic, horizontal lines that can be seen at regular intervals from the pleural line (A-lines, *Figure 1*, [Supplementary data online, Video S1](#)).⁷ When the air content in the lung decreases and lung density increases, vertical reverberation artefacts appear (B-lines, *Figure 2*, [Supplementary data online, Video S2](#)). B-lines originate from the pleural line and move synchronously with respiration, are laser-like in shape and extend towards the bottom of the US sector as displayed on the screen.⁸ B-lines are present in patients with HF and pulmonary oedema⁹ where an increasing number of visualized B-lines are associated with a decreasing air/water content ratio.^{10–12}

B-lines are not specific for cardiogenic pulmonary oedema and can be detected in patients with non-cardiogenic pulmonary oedema, including those with end stage renal disease and in ALI/ARDS,¹³ but also in pulmonary fibrosis (interstitial lung disease)¹⁴ and interstitial pneumonia. Some sonographic characteristics can help differentiate these various causes of B-lines⁵ (*Table 1*).

Pleural effusion can be easily detected by LUS, placing the phased-array transducer on the surface of the chest wall in an intercostal space, and is displayed as an anechoic space above the diaphragm. Pleural effusion is advised to be sought at first in the dependent zones, i.e. lateral and posterior chest wall (e.g. posterior axillary line) at the level of costophrenic angles (*Figure 3* and [Supplementary data online, Video S3](#)), which also allows other causes of chest X-ray radio-opacity, such as consolidation, mass, or an elevated hemidiaphragm to be ruled out (*Figure 4* and [Supplementary data online, Video S4](#)). LUS is more sensitive than chest X-ray in detecting pleural effusions, when using computed tomography as the reference standard,^{15,16} and can determine the volume of pleural effusion and monitor its evolution. LUS can also provide information on its likely nature by differentiating between simple and complex effusions, and can help guide the optimal site for needle thoracentesis.

Other sonographic features of congestion

The pathophysiology of congestion in HF includes the different stages of haemodynamic, pulmonary, and systemic congestion,¹⁷ which can be all assessed by ultrasound. Traditional echocardiography provides indicators of haemodynamic congestion, including a dilated left atrium, a high E/e', elevated pulmonary artery systolic pressure, and dilated inferior vena cava (IVC). B-lines on LUS are a sign of pulmonary congestion (increased extravascular lung water) due to left-sided HF, independent of—yet closely related with haemodynamic congestion. Chronically elevated left-sided filling pressures eventually result in a rise in right atrial pressures and

IVC distension. An IVC smaller than 21 mm that collapses >50% during inspiration suggests normal right atrial pressures,¹⁸ although this measurement should be interpreted in the context of the overall patient's underlying pathophysiological and haemodynamic status and integrated with other echocardiographic findings; an increased IVC diameter can identify intravascular volume expansion prior to changes in symptoms or body weight, and predicts a high risk of rehospitalization for HF or death in patients with acute or chronic HF.^{19,20}

Using a high-frequency linear transducer, the internal jugular vein (IJV) diameter can be measured. When congestion is severe, IJV distensibility, provoked by Valsalva manoeuvre, is markedly reduced, indicating a poor prognosis.²¹ Congestion in other organs can be also assessed by ultrasound, such as the kidneys. A comprehensive review of these novel techniques and their potential clinical utility can be found elsewhere.²² Recently, the venous excess ultrasound (VExUS) score, including Doppler evaluation of the IVC, hepatic veins, portal vein, and renal venous flow, has been proposed to assess presence and severity of venous congestion.²³ Whereas systemic venous congestion (as assessed by IVC, IJV, hepatic veins, portal veins, and renal veins) can be present in both right- and left-sided HF, LUS B-lines indicate pulmonary congestion due to left-sided HF.

Other sonographic features of deaeration

When the pulmonary air content is completely dissipated in areas of pulmonary consolidation, the lung parenchyma can be directly visualized (*Figure 5* and [Supplementary data online, Video S5](#)) with a hypoechoic or tissue-like pattern. Different aetiologies of consolidations may have also different sonographic appearance (*Figure 5*). Compression atelectasis is frequent in patients with HF; here, a large cardiogenic pleural effusion causes direct compression of the pulmonary parenchyma.⁴

On occasion, a focal interstitial syndrome (i.e. multiple B-lines localized only in a single area of the chest) can suggest pneumonia. These are thought to represent either the very early phases of pneumonia with the partial deaeration LUS pattern (B-lines) preceding the total deaeration LUS pattern (consolidation), or the focal oedema surrounding any consolidation.

LUS in HF: the technique

Whenever possible, it is advised to perform LUS exams with a standardized approach, to facilitate interpretation and monitoring, as well as reproducibility.²⁴ Several LUS protocols have been described, with a




Figure 1 The sonographic pattern of a normally aerated lung with the pleural line (upper dotted yellow line) and A-lines (lower dotted orange lines).




Figure 2 B-lines indicated by dotted white lines.

variable number of examination areas of the chest ('zones') to be examined, ranging from 4 to 28^{25–27} (Figure 6). Currently, the eight-zone scanning protocol is the most widely used, balancing the need for a simplified, rapid protocol, and good accuracy,⁸ with studies suggesting it the most appropriate approach across multiple settings, including the diagnosis of HF (with the 8-zone C-index non-inferior to the 28-zone C-index),²⁸ and for risk stratification.²⁹

When integrating LUS during stress echocardiography, a simplified four-zone scanning protocol is often used, placing the probe at the third intercostal space along the anterior axillary and mid-axillary lines.^{30,31}

In adult patients, both phased-array or convex transducers are sufficient to assess B-lines. If available, the abdominal preset (for the convex transducer) and the cardiac preset (for the phased-array transducer) usually provide adequate image quality. Some machines have also a dedicated lung preset. The transducer should be placed in the intercostal space either perpendicular (longitudinal and sagittal) or in parallel orientation (transverse) to the ribs. Imaging depth depends on the size of the patient, but is usually set at ~15–18 cm.²² Once the transducer position and gain settings are optimized for visualization of the pleural line and B-lines, it is advised to keep the transducer in the same location for at least one respiratory cycle. When a movie clip is recorded, the suggested length is of ~6 s.^{32,33} In each chest zone, the operator should scan all accessible chest surface to increase sensitivity.

During the examination, patients can be positioned either sitting upright, semi-recumbent, or supine; however, it is preferred to scan patients in the same position if serial examinations are being performed.³⁴ In those thoracic zones where a pleural effusion is visualized, B-lines cannot be assessed, and the presence of pleural effusions should be described in the LUS report instead.

There are two main approaches to quantify B-lines: score- or count-based methods. Score-based methods consider a minimum number of B-lines in one thoracic zone as a 'positive' zone (typically at least three B-lines),⁸ then, the number of positive zones are added up.²⁵ Count-based methods entail that B-lines are counted to obtain a number for each thoracic zone: B-lines can be counted one by one^{27,34} or, when confluent and overlapping, their number can be estimated from the percentage of space they occupy on the screen below the pleural line, divided by 10 (i.e. if ~60% of the screen below the pleural line is occupied by B-lines, it would conventionally count as six B-lines, up to a maximum of 10 per zone) (Figure 7). It is advised to count B-lines in the worst (less aerated/with more B-lines) point of each thoracic zone, then, the number of B-lines in each zone can be summed up

to obtain a total B-line count.^{4,8,35,36} All these methods have demonstrated good intra- and inter-observer agreement.^{37,38}

Key points: To evaluate pulmonary congestion in patients with suspected or established HF, using a phased-array or convex probe may be appropriate (including handheld ultrasound devices), in sagittal or transverse orientations (transducer perpendicular or parallel to the intercostal space), ideally following the eight-zone protocol and maintaining the patient in the same position during serial examinations.

Clinical applications

Acute heart failure

Diagnosis: ruling in/out acute cardiogenic dyspnoea

The typical LUS pattern of a patient with acute HF and pulmonary oedema is the presence of 'multiple' (at least three B-lines in one chest zone), 'diffuse' (at least two positive zones per hemithorax), and 'bilateral' B-lines. It is important to distinguish this pattern from the presence of a few patchy B-lines, especially when located at the lung bases, that can be seen even in healthy subjects, and which therefore do not qualify as 'multiple', 'diffuse', and 'bilateral'.

LUS is a rapid diagnostic test, which takes <5 min. Among patients hospitalized with acute dyspnoea, LUS has demonstrated high diagnostic accuracy for acute HF, with a sensitivity of 94–97% and a specificity of 97%^{25,26} that has been shown to be superior to both clinical assessment (i.e. history, physical examination, electrocardiogram, and arterial blood gas) and the chest X-ray, even when there is superimposed pneumonia.^{25,39,40} Indeed, LUS is superior to chest X-ray alone (sensitivity of 91.8% vs. 76.5% and a specificity of 92.3% vs. 87.0%),⁴¹ and has higher accuracy than clinical assessment even when integrated with 'traditional' diagnostic tests (i.e. natriuretic peptides and chest X-ray) to diagnose acute HF.^{25,26}

Monitoring: assessing decongestion

A higher number of B-lines on admission for an episode of acute HF is associated with greater risk, regardless of left ventricular ejection fraction (LVEF) and presence of HF signs and symptoms.⁴² This suggests that B-lines might be a potential target for treatment but also useful to monitor response to decongestive therapies.

Changes in B-lines are very dynamic and, for those who respond to diuretic therapy and other treatments, the number of B-lines decreases




Figure 3 The left costophrenic angle with pleural effusion.




Figure 4 Pleural effusion with compression atelectasis.

rapidly,^{27,43,44} regardless of the aetiology of acute HF.⁴⁵ Serial LUS can guide titration of diuretic therapy, leading to more rapid relief of congestion and, potentially, shortening the length of hospital stay.^{46–48} Resolution of pulmonary congestion detected by ultrasound during HF hospitalization is associated with improved survival (combined endpoint of 6 month all-cause death or acute heart failure rehospitalization).^{48,49} However, at the moment, there is no evidence that titrating diuretic therapy based on B-lines can improve morbidity and mortality in patients hospitalized with HF.⁴⁷

Although pleural effusions are typically monitored by chest X-ray in patients with acute HF, serial ultrasound scans are also effective in quantifying changes in the size of pleural effusion associated with decongestive therapy without exposing patients to radiation.⁵⁰

Prognosis: determining the timing of discharge

Up to 45% of patients hospitalized with acute HF are either readmitted for HF or die within 12 months of discharge.⁵¹ Those with a higher degree of residual congestion, including pulmonary congestion, at the time of discharge, are at particularly high risk.^{51,52} The importance of optimizing decongestion during a HF hospitalization is consequently highlighted in the 2021 ESC Guidelines.⁶ Patients with a high number of

B-lines at discharge are at higher risk of readmission or premature death than those discharged without or with only mild pulmonary congestion.^{35,36,52–54} Prior studies investigating the prevalence and prognostic value of B-lines at discharge in patients hospitalized with HF have employed 4, 8, or 28-zone LUS protocols; specific cut-off values for these protocols are summarized in Table 2. Collectively, these findings suggest that LUS may detect subclinical pulmonary congestion pre-discharge in patients with acute HF, irrespective of ejection fraction, and identify those at greater risk for subsequent adverse outcomes. In addition to B-lines, pleural effusion can be detected with ultrasound in approximately half of the patients discharged after an acute HF episode,²⁷ but it is unclear whether this finding is associated with an increased risk of cardiovascular events post-discharge.⁵⁶

Key points: The LUS pattern of pulmonary oedema in acute left-sided HF consists of the presence of multiple (at least three B-lines), diffuse (in at least two zones per hemithorax), and bilateral B-lines. Using LUS may be appropriate in patients with acute dyspnoea to rule in or rule out pulmonary oedema in suspected acute left-sided HF. Using LUS may be appropriate to detect persistent pulmonary congestion prior to discharge in patients hospitalized with acute HF, to identify those at higher risk of readmission for HF or death.




Figure 7 How to quantify B-lines.

LUS-guided strategy significantly improved the combined endpoint of urgent visit, rehospitalization, and death at 6 months after an acute HF episode.⁵⁸ Subsequently, the CLUSTER-HF trial also showed that LUS-guided treatment was associated with a 45% risk reduction in the primary endpoint (a composite of urgent HF visits, rehospitalization for worsening HF, and death from any cause), mainly driven by a reduction in urgent HF visits.⁵⁹ Another trial that randomized outpatients with chronic HF demonstrated a substantial reduction in hospitalizations for acute HF (by 56% at 90 days), as well as decreasing NT-proBNP concentrations and improving quality of life in those patients whose treatment was guided by LUS.⁶⁰ A summary of the clinical utility of LUS in acute and chronic heart failure is shown in Table 3.

Key points: Integrating LUS into the routine evaluation of ambulatory patients with HF may be appropriate to titrate diuretic therapy and reduce HF-related adverse outcomes, in particular, the need for urgent visits.

During stress echocardiography

LUS can be easily performed during exercise testing, including stress echocardiography.^{31,61,62} The number of B-lines increases during exercise in patients with HF more than in controls.^{63,64} In patients with heart failure with reduced ejection fraction (HFrEF), peak B-lines are closely associated with resting NT-proBNP, peak VO_2 , and stress pulmonary artery systolic pressure.⁶¹ In patients with heart failure with preserved ejection fraction (HFpEF), the development of pulmonary

congestion as measured by B-lines upon exercise correlates with exercise-induced worsening of diastolic function (particularly with peak E/e' and global strain rate during late diastole).⁶⁵ Haemodynamic studies further suggest that the acute development of pulmonary congestion during stress is related to an increase in pulmonary capillary hydrostatic pressure and systemic venous hypertension.⁶⁶ Therefore, LUS during exercise can quantify the pulmonary congestion secondary to the raise in stress-induced intracardiac/pulmonary pressures.

Detecting a higher number of B-lines at peak exercise is associated with a greater risk of HF hospitalization and death in patients with both HFrEF⁶¹ and HFpEF. Overall, these findings suggest that LUS may complement standard exercise testing to improve HF diagnosis and risk stratification.^{67,68}

In patients with chest pain and acute myocardial infarction

Patients with acute coronary syndromes may develop pulmonary congestion, especially during or soon after an ST-elevation myocardial infarction. A recent systematic review suggests that a high number of B-lines is a common finding in patients hospitalized with an acute coronary syndrome and identifies patients more likely to have adverse in-hospital and long-term outcomes.⁶⁹ Whether assessment of B-lines might guide use of therapies that reduce pulmonary congestion after a myocardial infarction and improve morbidity and mortality in the short-term and long-term remains to be determined.

Table 2 Overview of select pre-discharge imaging protocols and cut-off values in acute HF

Number of zones	B-line cut-off	Ultrasound equipment	HF readmission or death	Reference
4	≥7 B-lines	High-end system; phased-array	90 days: Adj. HR 3.03 (95% CI 1.45 to 6.31)	Platz 2019 ²⁷
8	≥1 zone with ≥3 B-lines (one positive zone) on each hemithorax	High-end system; phased-array	90 days: Adj. HR 3.30 (95% CI 1.00 to 10.91)	Coiro 2015 ³⁵
28	>15 B-lines	High-end system; phased-array	180 days: Adj. HR 11.74 (95% CI 1.30 to 106.16)	Gargani 2015 ³⁶

Adj., adjusted; CI, confidence interval; HF, heart failure; HR, hazard ratio.

Table 3 LUS in acute and chronic heart failure

	Aim	LUS picture	Advantages
Acute heart failure			
Diagnosis	Rule in and rule out acute left-sided HF in patients with dyspnoea.	Multiple, diffuse, bilateral B-lines rule in AHF. Absence of multiple, diffuse, bilateral B-lines rules out AHF.	LUS improves diagnostic accuracy compared with standard strategy (chest X-ray + NT-proBNP). LUS reduces time to correct diagnosis. LUS detects subclinical pulmonary congestion in patients with HF who have mild or absent signs and symptoms.
Monitoring	Monitor decongestion during AHF hospitalization.	Reduction of the number of B-lines. Reduction of the size of pleural effusion, if any.	Bedside monitoring to support decision-making in diuretic therapy and fluid management. A significant reduction of B-lines during hospitalization is associated with a lower risk of rehospitalization for AHF and death at 6 months.
Prognosis	Determine the timing of discharge. Detect persistent subclinical congestion at discharge to identify patients at higher risk of rehospitalization for AHF or death.	Persistent B-lines at discharge, even with resolved signs and symptoms of HF.	A high number of B-lines at discharge predict rehospitalization for AHF and death at 3 months.
Chronic heart failure			
During follow-up	Guide diuretic therapy after discharge.	Reduction of the number of B-lines. Reduction of the size of pleural effusion, if any.	LUS-guided therapy might reduce risk of urgent HF visits and hospital admission for AHF.
Prognosis	Identification of patients at higher risk of rehospitalization for AHF or death.	Presence of B-lines at out-office visits, even without signs and symptoms of HF.	An elevated number of B-lines in outpatients predict hospitalization for AHF and death at 6 months.

LUS, lung ultrasound; HF, heart failure; AHF, acute heart failure.





In patients with shock/hypotension

LUS can provide vital information when integrated with echocardiography in the assessment of hypotension and various types of shock, with the potential of detecting the underlying cause, and improving the quality of care delivered to critically ill patients.

In hypovolaemic shock, the presence of a normal LUS pattern confirms the presence of ‘dry’ lungs that has a very high negative predictive value in ruling out a cardiogenic cause of hypotension^{2,70} (Figure 8). Patients in cardiogenic shock due to LV failure or mitral/aortic valvular heart disease eventually develop pulmonary congestion that would

appear by LUS as multiple, diffuse, and bilateral B-lines, associated with a regular pleural line¹³ (Figure 8).

In patients with a pneumothorax, LUS has superior diagnostic sensitivity (88% vs. 52%) and similar specificity (99% vs. 100%) to chest X-ray.⁷¹ The combination of the absence of lung sliding (generating the barcode sign on M-mode), absence of B-lines, and the presence of a lung point (the transition point between a LUS pattern with lung sliding and without lung sliding) would suggest a pneumothorax as the cause of hypotension/shock (Figure 8).⁸ The visualization of a normal LUS pattern along with non-compressible peripheral deep veins due to a thrombus would suggest pulmonary embolism, especially when coupled with RV

Table 4 How to integrate echocardiographic and pulmonary findings to assess congestion status		
E/e' and other echocardiographic signs of increased left ventricular filling pressures		B-lines
Normal		No B-lines No congestion
Increased		No B-lines Haemodynamic congestion
Increased		Multiple diffuse bilateral B-lines Haemodynamic and pulmonary congestion
Normal		Multiple diffuse bilateral B-lines Pulmonary congestion without haemodynamic congestion (check for ALI/ARDS or other causes of B-lines)

ALI, acute lung injury; ARDS, acute respiratory distress syndrome.

dilatation and dysfunction.⁷² Moreover, triangular/polygonal hypoechoic peripheral consolidations with sharp margins may be detected, as the sonographic appearance of pulmonary infarctions (Figure 8).⁷³ In septic shock, the lungs are a frequent source of sepsis, due to lobar pneumonia that can be diagnosed with LUS with high sensitivity, through the visualization of consolidations often with dynamic

air bronchograms, commonly at the most dependant zones of the lungs, with or without adjacent pleural effusion. The presence of punctiform echoes within an otherwise anechogenic pleural effusion could indicate exudative pleural effusion/empyema.⁷⁴ The lungs can develop B-lines due to ARDS because of septic shock of extrapulmonary origin. LUS can be very useful in monitoring areation/




Figure 8 Integrated cardiopulmonary ultrasound for the differential diagnosis of respiratory and circulatory failure.

deareation, assessing disease progression and recruitment in ALI/ARDS (including COVID-19 pneumonia).⁷⁵ LUS can be performed also during transoesophageal echocardiography through dedicated views of the posterior regions of the lungs where lung consolidations and pleural effusions are most often seen, with promising preliminary results.⁷⁶

Key points: In patients with shock/hypotension, LUS can complement the standard clinical evaluation to improve diagnostic accuracy.

Integrated cardiopulmonary ultrasound

How to integrate echocardiography and LUS

Transthoracic echocardiography is guidelines-recommended as a key investigation in HF⁶ for the assessment of cardiac structure and function, defining HF phenotype based on LVEF measurement, as well identifying potential underlying aetiologies and mechanisms, and guiding the introduction of treatment. The 2021 ESC Guidelines also mention the use of LUS in acute settings to confirm a diagnosis of HF, especially when natriuretic peptide testing is not available,⁶ since pulmonary congestion is one of the main causes of hospital admission in patients with HF.¹⁷

Integrating cardiac and pulmonary assessments on ultrasound can help understand and identify the pulmonary consequences of structural or functional cardiac pathology, and potentially individualize patient management. Real-time interpretation of B-lines, together with the non-invasive haemodynamic signs that can be measured by echocardiography, can provide simultaneous assessments of the cardiac and pulmonary state, leading to more accurate diagnosis and earlier treatment initiation.^{77,78} The number of B-lines is related to invasive left ventricular end-diastolic pressure measurements,⁷⁹ to E/e',⁸⁰ to tricuspid

regurgitation velocity,⁴² to right ventricular (RV) function, and RV–pulmonary artery coupling.⁸¹ However, in patients with the same haemodynamic profile, the degree of pulmonary congestion as detected by B-lines can vary markedly. Therefore, echocardiography is useful to understand the aetiology of HF, whereas LUS B-lines are especially useful to detect the degree of pulmonary congestion. The routine assessment of B-lines may be particularly appropriate when either a high-end ultrasound machine or the expertise for more complex echocardiographic analyses is not readily available or when cardiac image quality is poor and initial diagnosis difficult. Therefore, whenever possible, these two aspects of the patient's compensation status—haemodynamic congestion and pulmonary congestion—¹⁷ can be evaluated in an integrated ultrasound cardiopulmonary examination (Table 4). This approach can also provide concomitant information regarding perfusion (by the LV outflow tract time-velocity integral as a non-invasive index of cardiac output) and pulmonary congestion (by B-lines), in a less invasive way compared with right heart catheterization. The integration of B-lines with IVC is also valuable and integrates different information, since IVC diameter and collapsibility index are proxies for right atrial pressure, but also reflects intravascular volume status, whereas B-lines reflect extravascular lung water, which can be present in patients with either a normal or a dilated IVC.¹⁹

Key points: When evaluating patients with known or suspected HF, it is advised to integrate LUS B-lines in the echocardiographic assessment to provide information about the degree of pulmonary congestion, in both acute and chronic settings.

LUS in FoCUS

Focus cardiac ultrasound (FoCUS) is a point-of-care cardiac ultrasound examination performed according to a standardized, but restricted, scanning protocol in patients presenting with circulatory or respiratory compromise, chest pain or trauma, dyspnoea, syncope, or cardiac arrest.⁸² The FoCUS scanning protocol comprises five echocardiographic

Setting	Trials/studies needed		Main questions to be addressed
Acute heart failure	Additional randomized trials assessing whether LUS can improve the diagnosis and characterization of patients with AHF beyond traditional approaches.	• Can the use of LUS facilitate the classification of AHF phenotypes that may benefit from specific therapeutic interventions? • Can LUS imaging be used to improve management decisions in patients with AHF, e.g. hospital admission vs. early discharge from the Emergency Department? • Is the use of LUS imaging effective and safe as part of a strategy to improve congestion relief in AHF, including intensification/de-intensification of diuretic therapy? • Can the use of LUS improve post-discharge outcomes in patients with AHF?	
Chronic heart failure	Development and validation of diagnostic protocols assessing the utility of LUS at rest or during exercise in the diagnosis of HFpEF.	• Can the use of LUS improve the clinical course of patients with chronic HF (HFrEF and/or HFpEF)? • Can the use of LUS facilitate the identification of subsets of patients with chronic HF that can benefit from specific therapeutic interventions? • Is the use of LUS in the management of patients with chronic HF cost-effective? • Can the use of LUS improve the clinical course of specific HF phenotypes (myocarditis, cardiotoxicity, inherited cardiomyopathies, post-partum cardiomyopathy, and cardiac amyloidosis)?	
Acute coronary syndromes	Additional randomized trials assessing whether LUS can improve the diagnosis of the aetiology of chest pain beyond traditional approaches.	• Can the use of LUS facilitate the identification of subsets of patients with ACS that may benefit from specific therapeutic interventions?	

AHF, acute heart failure; HF, heart failure; HFrEF, heart failure with reduced ejection fraction; HFpEF, heart failure with preserved ejection fraction; LUS, lung ultrasound.

views sufficient for crude assessment of left and right ventricular size and function, pericardial effusion, and intravascular volume status.

Since cardiovascular and pulmonary diseases frequently coexist and may share clinical presentations, LUS may help in the differential diagnosis of acute dyspnoea and haemodynamic instability and is therefore an integral part of the FoCUS examination.⁸² The presence of multiple, diffuse, and bilateral B-lines in a dyspnoeic patient with LV dysfunction is almost unequivocally consistent with HFrEF. In the presence of left atrial dilation and/or visually-detected LV hypertrophy, multiple, diffuse, and bilateral B-lines may indicate HFpEF, and natriuretic peptide measurement and a comprehensive echocardiography should be ordered for further evaluation.⁸² Figure 8 summarizes the main situations of respiratory and circulatory failure, where LUS integrated with FoCUS can be of particular value.

Key points: Whenever possible, it is advised to integrate assessment of B-lines and pleural effusion into the FoCUS examination since it may help in the differential diagnosis of acute dyspnoea and haemodynamic instability.

Pitfalls

As with every diagnostic technique, there are some limitations with LUS examinations, which should be acknowledged to master this tool. There are still uncertainties about the exact physical origin of B-lines,¹¹ although they are clearly related to a partial deaeration of the pulmonary parenchyma. LUS is an easy to learn ultrasound technique, with a short learning curve that can be limited to about 20 exams,^{83,84} but some training is needed to ensure reproducibility and the correct interpretation of findings. Initiatives that would ensure quality standards and safe patient care, for instance, a LUS certification programme, could facilitate the standardization of this procedure for both clinicians and other healthcare professionals involved in the care of patients with HF, who can successfully assess B-lines.^{85,86} B-lines are artefacts: although no clinically significant differences have been reported by using different machines and settings, attention should be paid to avoid magnification or deletion of artefactual images. Different LUS scanning

protocols have been proposed and used, in research and in clinical practice, as previously discussed.

Probably, the main pitfall is the clinical interpretation of B-lines in the absence of an established diagnosis. B-lines are not specific for cardiogenic pulmonary oedema: being a sign of partial pulmonary deaeration, they can be present in patients with interstitial lung disease, such as pulmonary fibrosis, as well as ALI/ARDS, and interstitial pneumonia (including COVID-19 pneumonia). As highlighted in Table 1, there are some LUS features that can help differentiate these conditions, taking into account that a strong integration with clinical and other routine diagnostic findings is mandatory. This point further reinforces the need for LUS training, to ensure that healthcare providers will not categorize every patient with B-lines as having HF.

Importantly, the absence of substantial pulmonary congestion on ultrasound does not exclude HF, particularly in the context of severe obesity or isolated right-sided HF; therefore, clinicians should not forget to evaluate systemic/venous congestion.⁸⁷

Gaps in current evidence

In recent years, major advances have been made in our understanding of the utility of LUS as an imaging tool in the diagnosis, monitoring, and risk stratification of patients with HF, but areas with lack of evidence remain. Table 5 lists selected topics that deserve to be addressed in future clinical research.

While waiting for additional evidence regarding the clinical utility of LUS, especially in titrating diuretic therapy, we already have solid data supporting the use of this tool in daily clinical practice to complement either FoCUS or comprehensive echocardiography in the assessment and management of patients with HF.

Conclusions

LUS is an easy to learn, rapid, and versatile point-of-care diagnostic method. B-lines on LUS are the sonographic sign of a partially deaerated

lung, which can be detected in patients with HF, providing a semi-quantitative evaluation of pulmonary interstitial oedema.

Sonographic B-lines are useful for the diagnosis, monitoring, and prognostic assessment of patients with HF. Therefore, it would be important to integrate routine evaluation of B-lines and pleural effusions into standard transthoracic echocardiographic protocols in patients with suspected or confirmed HF, as well as in the FoCUS examination. This integrated cardiopulmonary ultrasound approach has the potential to improve clinical management and outcomes, but further research is required.

Supplementary data

Supplementary data is available at *European Heart Journal - Cardiovascular Imaging* online.

Funding

None declared.

Conflict of interest: Dr. Gargani: no conflict of interest related to this paper. Outside of this paper, Dr. Gargani has received consultancy honoraria from Philips Healthcare, GE Healthcare, Caption Health, EchoNous. Dr. Girerd: no conflict of interest related to this paper. Outside of this paper, Dr. Girerd has received consultancy honoraria from Astra Zeneca, Bayer, Boehringer-Ingelheim, Novartis, Lilly, Vifor International. Dr. Platz: no conflict of interest related to this paper. Outside of this paper, Dr. Platz' employer has received support from Novartis for consulting work, and she has consulted for scPharmaceuticals. She has received research support from the NIH (R01HL148439), the American Heart Association and AstraZeneca. Dr. Pellicori: no conflict of interest related to this paper. Outside of this paper, Dr. Pellicori has received consultancy honoraria and/or sponsorship support from Boehringer Ingelheim, Pharmacosmos, Novartis, Vifor, AstraZeneca and Caption Health, and research support from Bristol Myers Squibb. Dr. Stankovic: no conflict of interest related to this paper. Outside of this paper, Dr. Stankovic has received consultancy honoraria from Servier, Boehringer-Ingelheim, Pfizer, Astra Zeneca, Novartis, Merck Sharp & Dohme. Dr. Palazzuoli: no conflict of interest to declare. Dr. Pivetta: no conflict of interest related to this paper. Outside of this paper, Dr. Pivetta has received research support from Butterfly inc. Dr. Miglioranza: no conflict of interest related to this paper. Outside of this paper, Dr. Miglioranza has received consultancy honoraria from Philips, GE Healthcare, Abbott. Dr. Aboumarie: no conflict of interest to declare. Dr. Agricola: no conflict of interest related to this paper. Outside of this paper, Dr. Agricola has received consultancy honoraria from Philips, Edwards Lifesciences. Dr. Volpicelli: no conflict of interest to declare. Dr. Price: no conflict of interest to declare. Dr. Donal: no conflict of interest related to this paper. Outside of this paper, Dr. Donal has received research funding from Abbott. Dr. Cosyns: no conflict of interest related to this paper. Outside of this paper, Dr. Cosyns has received research funding from Abbott. Dr. Neskovic: no conflict of interest related to this paper. Outside of this paper, Dr. Neskovic has received consultancy honoraria from Bayer, AstraZeneca, Pfizer, Boehringer Ingelheim, Servier and Novartis.

Data availability

No new data were generated or analysed in support of this research.

References

- Lichtenstein D, Mezière G. A lung ultrasound sign allowing bedside distinction between pulmonary edema and COPD: the comet-tail artifact. *Intensive Care Med* 1998;**24**: 1331–4.
- Lichtenstein DA, Mezière GA. Relevance of lung ultrasound in the diagnosis of acute respiratory failure: the BLUE protocol. *Chest* 2008;**134**:117–25.
- Picano E, Frassi F, Agricola E, Gligorova S, Gargani L, Mottola G. Ultrasound lung comets: a clinically useful sign of extravascular lung water. *J Am Soc Echocardiogr* 2006;**19**:356–63.
- Gargani L. Ultrasound of the lungs: more than a room with a view. *Heart Fail Clin* 2019;**15**:297–303.
- Gargani L. Lung ultrasound: a new tool for the cardiologist. *Cardiovasc Ultrasound* 2011;**9**:6.
- McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M et al. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;**42**:3599–726.
- Gargani L, Volpicelli G. How I do it: lung ultrasound. *Cardiovasc Ultrasound* 2014;**12**: 1–10.
- Volpicelli G, Elbarbary M, Blaivas M, Lichtenstein DA, Mathis G, Kirkpatrick AW et al. International evidence-based recommendations for point-of-care lung ultrasound. *Intensive Care Med* 2012;**38**:577–91.
- Lichtenstein D, Mezière G, Biderman P, Gepner A, Barré O. The comet-tail artifact. An ultrasound sign of alveolar-interstitial syndrome. *Am J Respir Crit Care Med* 1997;**156**: 1640–6.
- Jambrik Z, Gargani L, Adamicz A, Kaszaki J, Varga A, Forster T et al. B-lines quantify the lung water content: a lung ultrasound versus lung gravimetry study in acute lung injury. *Ultrasound Med Biol* 2010;**36**:2004–10.
- Soldati G, Inchingolo R, Smargiassi A, Sher S, Nenna R, Inchingolo CD et al. Ex vivo lung sonography: morphologic-ultrasound relationship. *Ultrasound Med Biol* 2012;**38**: 1169–79.
- Gargani L, Lionetti V, Di Cristofano C, Bevilacqua G, Recchia FA, Picano E. Early detection of acute lung injury uncoupled to hypoxemia in pigs using ultrasound lung comets. *Crit Care Med* 2007;**35**:2769–74.
- Copetti R, Soldati G, Copetti P. Chest sonography: a useful tool to differentiate acute cardiogenic pulmonary edema from acute respiratory distress syndrome. *Cardiovasc Ultrasound* 2008;**6**:16.
- Gargani L, Bruni C, Romei C, Frumento P, Moreo A, Agoston G et al. Prognostic value of lung ultrasound B-lines in systemic sclerosis. *Chest* 2020;**158**:1515–25.
- Balik M, Plasil P, Waldauf P, Pazout J, Fric M, Otahal M et al. Ultrasound estimation of volume of pleural fluid in mechanically ventilated patients. *Intensive Care Med* 2006;**32**:318–21.
- Roch A, Bojan M, Michelet P, Romain F, Bregeon F, Papazian L et al. Usefulness of ultrasonography in predicting pleural effusions >500 mL in patients receiving mechanical ventilation. *Chest* 2005;**127**:224–32.
- Gheorghiade M, Follath F, Ponikowski P, Barsuk JH, Blair JE, Cleland JG et al. Assessing and grading congestion in acute heart failure: a scientific statement from the acute heart failure committee of the heart failure association of the European Society of Cardiology and endorsed by the European Society of Intensive Care Medicine. *Eur J Heart Fail* 2010;**12**:423–33.
- Rudski LG, Lai WW, Afilalo J, Hua L, Handschumacher MD, Chandrasekaran K et al. Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology, and the Canadian Society of Echocardiography. *J Am Soc Echocardiogr* 2010;**23**:685–713.
- Pellicori P, Carubelli V, Zhang J, Castiello T, Sherwi N, Clark AL et al. IVC diameter in patients with chronic heart failure: relationships and prognostic significance. *JACC Cardiovasc Imaging* 2013;**6**:16–28.
- Pellicori P, Shah P, Cuthbert J, Urbinati A, Zhang J, Kallvikbacka-Bennett A et al. Prevalence, pattern and clinical relevance of ultrasound indices of congestion in outpatients with heart failure. *Eur J Heart Fail* 2019;**21**:904–16.
- Pellicori P, Kallvikbacka-Bennett A, Dierckx R, Zhang J, Putzu P, Cuthbert J et al. Prognostic significance of ultrasound-assessed jugular vein distensibility in heart failure. *Heart* 2015;**101**:1149–58.
- Pellicori P, Platz E, Dauw J, Ter Maaten JM, Martens P, Pivetta E et al. Ultrasound imaging of congestion in heart failure: examinations beyond the heart. *Eur J Heart Fail* 2021;**23**: 703–12.
- Beaubien-Souligny W, Rola P, Haycock K, Bouchard J, Lamarche Y, Spiegel R et al. Quantifying systemic congestion with point-of-care ultrasound: development of the venous excess ultrasound grading system. *Ultrasound J* 2020;**12**:16.
- Platz E, Jhund PS, Girerd N, Pivetta E, McMurray JJ V, Peacock WF et al. Expert consensus document: reporting checklist for quantification of pulmonary congestion by lung ultrasound in heart failure. *Eur J Heart Fail* 2019;**21**:844–51.
- Pivetta E, Goffi A, Lupia E, Tizzani M, Porrino G, Ferreri E et al. Lung ultrasound-implemented diagnosis of acute decompensated heart failure in the emergency department—a SIMEU multicenter study. *Chest* 2015;**148**:202–10.
- Pivetta E, Goffi A, Nazerian P, Castagno D, Tozzetti C, Tizzani P et al. Lung ultrasound integrated with clinical assessment for the diagnosis of acute decompensated heart failure in the emergency department: a randomized controlled trial. *Eur J Heart Fail* 2019;**21**:754–66.

27. Platz E, Campbell RT, Claggett B, Lewis EF, Groarke JD, Docherty KF *et al.* Lung ultrasound in acute heart failure: prevalence of pulmonary congestion and short- and long-term outcomes. *JACC Heart Fail* 2019;**7**:849–58.
28. Buessler A, Chouhied T, Duarte K, Bassand A, Huot-Marchand M, Gottwalles Y *et al.* Accuracy of several lung ultrasound methods for the diagnosis of acute heart failure in the ED: a multicenter prospective study. *Chest* 2020;**157**:99–110.
29. Rastogi T, Bozec E, Pellicori P, Bayes-Genis A, Coiro S, Domingo M *et al.* Prognostic value and therapeutic utility of lung ultrasound in acute and chronic heart failure: a meta-analysis. *JACC Cardiovasc Imaging* 2022;**15**:950–2.
30. Scali MC, Zagatina A, Simova I, Zhuravskaya N, Ciampi Q, Paterni M *et al.* B-lines with lung ultrasound: the optimal scan technique at rest and during stress. *Ultrasound Med Biol* 2017;**43**:2558–66.
31. Merli E, Ciampi Q, Scali MC, Zagatina A, Merlo PM, Arbucci R *et al.* Pulmonary congestion during exercise stress echocardiography in ischemic and heart failure patients. *Circ Cardiovasc Imaging* 2022;**15**:e013558.
32. Pivetta E, Baldassa F, Masellis S, Bovo F, Lupia E, Maule MM. Sources of variability in the detection of B-lines, using lung ultrasound. *Ultrasound Med Biol* 2018;**44**:1212–6.
33. Platz E, Pivetta E, Merz AA, Peck J, Rivero J, Cheng S. Impact of device selection and clip duration on lung ultrasound assessment in patients with heart failure. *Am J Emerg Med* 2015;**33**:1552–6.
34. Frasure SE, Matlisky DK, Siadecki SD, Platz E, Saul T, Lewis RE. Impact of patient positioning on lung ultrasound findings in acute heart failure. *Eur Heart J Acute Cardiovasc Care* 2015;**4**:326–32.
35. Coiro S, Rossignol P, Ambrosio G, Carluccio E, Alunni G, Murrone A *et al.* Prognostic value of residual pulmonary congestion at discharge assessed by lung ultrasound imaging in heart failure. *Eur J Heart Fail* 2015;**17**:1172–81.
36. Gargani L, Pang PSS, Frassi F, Miglioranza MH, Dini FLL, Landi P *et al.* Persistent pulmonary congestion before discharge predicts rehospitalization in heart failure: a lung ultrasound study. *Cardiovasc Ultrasound* 2015;**13**:40.
37. Jambrik Z, Monti S, Coppola V, Agricola E, Mottola G, Miniati M *et al.* Usefulness of ultrasound lung comets as a nonradiologic sign of extravascular lung water. *Am J Cardiol* 2004;**93**:1265–70.
38. Gargani L, Sicari R, Raciti M, Serasini L, Passera M, Torino C *et al.* Efficacy of a remote web-based lung ultrasound training for nephrologists and cardiologists: a LUST trial sub-project. *Nephrol Dial Transplant* 2016;**31**:1982–8.
39. Cibinel GA, Casoli G, Elia F, Padoan M, Pivetta E, Lupia E *et al.* Diagnostic accuracy and reproducibility of pleural and lung ultrasound in discriminating cardiogenic causes of acute dyspnea in the emergency department. *Intern Emerg Med* 2012;**7**:65–70.
40. Mazzola M, Pugliese NR, Zavagli M, De Biase N, Bandini G, Barbarisi G *et al.* Diagnostic and prognostic value of lung ultrasound B-lines in acute heart failure with concomitant pneumonia. *Front Cardiovasc Med* 2021;**8**:693912.
41. Chiu L, Jairam MP, Chow R, Chiu N, Shen M, Alhassan A *et al.* Meta-analysis of point-of-care lung ultrasonography versus chest radiography in adults with symptoms of acute decompensated heart failure. *Am J Cardiol* 2022;**174**:89–95.
42. Gargani L, Pugliese NR, Frassi F, Frumento P, Poggiani E, Mazzola M *et al.* Prognostic value of lung ultrasound in patients hospitalized for heart disease irrespective of symptoms and ejection fraction. *ESC Heart Fail* 2021;**8**:2660–9.
43. Volpicelli G, Caramello V, Cardinale L, MUSA A, Bar F, Frascisco MF. Bedside ultrasound of the lung for the monitoring of acute decompensated heart failure. *Am J Emerg Med* 2008;**26**:585–91.
44. Cortellaro F, Ceriani E, Spinelli M, Campanella C, Bossi I, Coen D *et al.* Lung ultrasound for monitoring cardiogenic pulmonary edema. *Intern Emerg Med* 2016;**12**:1011–7.
45. Martindale JL, Secko M, Kilpatrick JF, Ian S, Paladino L, Aherne A *et al.* Serial sonographic assessment of pulmonary edema in patients with hypertensive acute heart failure. *J Ultrasound Med* 2017;**37**:337–45.
46. Mozzini C, Di Dio Perna M, Perna D, Pesce G, Garbin U, Meschi T *et al.* Lung ultrasound in internal medicine efficiently drives the management of patients with heart failure and speeds up the discharge time inferior cave vein collapsibility index. *Intern Emerg Med* 2017;**13**:27–33.
47. Pang PS, Russell FM, Ehrman R, Ferre R, Gargani L, Levy PD *et al.* Lung Ultrasound-Guided Emergency Department Management of Acute Heart Failure (BLUSHED-AHF): a randomized controlled pilot trial. *JACC Heart Fail* 2021;**9**:638–48.
48. Öhman J, Harjola VP, Karjalainen P, Lassus J. Focused echocardiography and lung ultrasound protocol for guiding treatment in acute heart failure. *ESC Heart Fail* 2018;**5**:120–8.
49. Öhman J, Harjola VP, Karjalainen P, Lassus J. Assessment of early treatment response by rapid cardiothoracic ultrasound in acute heart failure: cardiac filling pressures, pulmonary congestion and mortality. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:311–20.
50. Brogi E, Gargani L, Bignami E, Barbariol F, Marra A, Forfori F *et al.* Thoracic ultrasound for pleural effusion in the intensive care unit: a narrative review from diagnosis to treatment. *Critical Care* 2017;**21**:325.
51. Ambrosy AP, Pang PS, Khan S, Konstam MA, Fonarow GC, Traver B *et al.* Clinical course and predictive value of congestion during hospitalization in patients admitted for worsening signs and symptoms of heart failure with reduced ejection fraction: findings from the EVEREST trial. *Eur Heart J* 2013;**34**:835–43.
52. Chioncel O, Mebazaa A, Maggioni AP, Harjola VP, Rosano G, Laroche C *et al.* Acute heart failure congestion and perfusion status—impact of the clinical classification on in-hospital and long-term outcomes: insights from the ESC-EORP-HFA Heart Failure Long-Term Registry. *Eur J Heart Fail* 2019;**21**:1338–52.
53. Cogliati C, Casazza G, Ceriani E, Torzillo D, Furlotti S, Bossi I *et al.* Lung ultrasound and short-term prognosis in heart failure patients. *Int J Cardiol* 2016;**218**:104–8.
54. Palazzuoli A, Ruocco G, Beltrami M, Nuti R, Cleland JG. Combined use of lung ultrasound, B-type natriuretic peptide, and echocardiography for outcome prediction in patients with acute HFrEF and HFpEF. *Clin Res Cardiol* 2018;**107**:586–96.
55. Coiro S, Porot G, Rossignol P, Ambrosio G, Carluccio E, Tritto I *et al.* Prognostic value of pulmonary congestion assessed by lung ultrasound imaging during heart failure hospitalisation: a two-centre cohort study. *Sci Rep* 2016;**6**:39426.
56. Lindner M, Thomas R, Claggett B, Lewis EF, Groarke J, Merz AA *et al.* Quantification of pleural effusions on thoracic ultrasound in acute heart failure. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:513–21.
57. Pellicori P, Cleland JGF, Zhang J, Kallvikbacka-Bennett A, Urbinati A, Shah P *et al.* Cardiac dysfunction, congestion and loop diuretics: their relationship to prognosis in heart failure. *Cardiovasc Drugs Ther* 2016;**30**:599–609.
58. Rivas-Lasarte M, Álvarez-García J, Fernández-Martínez J, Maestro A, López-López L, Solé-González E *et al.* Lung ultrasound-guided treatment in ambulatory patients with heart failure: a randomized controlled clinical trial (LUS-HF study). *Eur J Heart Fail* 2019;**21**:1605–13.
59. Araiza-Garaygordobil D, Gopar-Nieto R, Martínez-Amezcuca P, Cabello-López A, Alanis-Estrada G, Luna-Herbert A *et al.* A randomized controlled trial of lung ultrasound-guided therapy in heart failure (CLUSTER-HF study). *Am Heart J* 2020;**227**:31–9.
60. Marini C, Fragasso G, Italia L, Sisakian H, Tufaro V, Ingallina G *et al.* Lung ultrasound-guided therapy reduces acute decompensation events in chronic heart failure. *Heart* 2020;**106**:1934–9.
61. Scali MC, Cortigiani L, Simionuc A, Gregori D, Marzilli M, Picano E. Exercise-induced B-lines identify worse functional and prognostic stage in heart failure patients with depressed left ventricular ejection fraction. *Eur J Heart Fail* 2017;**19**:1468–78.
62. Agricola E, Picano E, Oppizzi M, Pisani M, Meris A, Fragasso G *et al.* Assessment of stress-induced pulmonary interstitial edema by chest ultrasound during exercise echocardiography and its correlation with left ventricular function. *J Am Soc Echocardiogr* 2006;**19**:457–63.
63. Simonovic D, Coiro S, Carluccio E, Girerd N, Deljanin-Ilic M, Cattadori G *et al.* Exercise elicits dynamic changes in extravascular lung water and haemodynamic congestion in heart failure patients with preserved ejection fraction. *Eur J Heart Fail* 2018;**20**:1366–9.
64. Pugliese NR, Mazzola M, Fabiani I, Gargani L, De Biase N, Pedrinelli R *et al.* Haemodynamic and metabolic phenotyping of hypertensive patients with and without heart failure by combining cardiopulmonary and echocardiographic stress test. *Eur J Heart Fail* 2020;**22**:458–68.
65. Simonovic D, Coiro S, Deljanin-Ilic M, Kobayashi M, Carluccio E, Girerd N *et al.* Exercise-induced B-lines in heart failure with preserved ejection fraction occur along with diastolic function worsening. *ESC Heart Fail* 2021;**8**:5068–80.
66. Reddy YNV, Obokata M, Wiley B, Koepp KE, Jorgenson CC, Egbe A *et al.* The haemodynamic basis of lung congestion during exercise in heart failure with preserved ejection fraction. *Eur Heart J* 2019;**40**:3721–30.
67. Coiro S, Simonovic D, Deljanin-Ilic M, Duarte K, Carluccio E, Cattadori G *et al.* Prognostic value of dynamic changes in pulmonary congestion during exercise stress echocardiography in heart failure with preserved ejection fraction. *Circ Heart Fail* 2020;**13**:e006769.
68. Pugliese NR, De Biase N, Gargani L, Mazzola M, Conte L, Fabiani I *et al.* Predicting the transition to and progression of heart failure with preserved ejection fraction: a weighted risk score using bio-humoral, cardiopulmonary, and echocardiographic stress testing. *Eur J Prev Cardiol* 2021;**28**:1650–61.
69. Lindner M, Lindsey A, Bain PA, Platz E. Prevalence and prognostic importance of lung ultrasound findings in acute coronary syndrome: a systematic review. *Echocardiography* 2021;**38**:2069–76.
70. Lancellotti P, Price S, Edvardsen T, Cosyns B, Neskovic AN, Dulgheru R *et al.* The use of echocardiography in acute cardiovascular care: recommendations of the European Association of Cardiovascular Imaging and the Acute Cardiovascular Care Association. *Eur Heart J Cardiovasc Imaging* 2015;**16**:119–46.
71. Ding W, Shen Y, Yang J, He X, Zhang M. Diagnosis of pneumothorax by radiography and ultrasonography: a meta-analysis. *Chest* 2011;**140**:859–66.
72. Nazerian P, Vanni S, Volpicelli G, Gigli C, Zanobetti M, Bartolucci M *et al.* Accuracy of point-of-care multiorgan ultrasonography for the diagnosis of pulmonary embolism. *Chest* 2013;**145**:950–7.
73. Copetti R, Cominotto F, Meduri S, Orso D. The 'survived lung': an ultrasound sign of 'bubbly consolidation' pulmonary infarction. *Ultrasound Med Biol* 2020;**46**:2546–50.
74. Lichtenstein DA. Ultrasound in the management of thoracic disease. *Crit Care Med* 2007;**35**:S250–61.

75. Gargani L, Soliman-Aboumarie H, Volpicelli G, Corradi F, Pastore MC, Cameli M. Why, when, and how to use lung ultrasound during the COVID-19 pandemic: enthusiasm and caution. *Eur Heart J Cardiovasc Imaging* 2020;**21**:941–8.
76. Cavayas YA, Girard M, Desjardins G, Denault AY. Transesophageal lung ultrasonography: a novel technique for investigating hypoxemia. *Can J Anesthesia* 2016;**63**:1266–76.
77. Russell FM, Ehrman RR, Cosby K, Ansari A, Tseeng S, Christain E et al. Diagnosing acute heart failure in patients with undifferentiated dyspnea: a lung and cardiac ultrasound (LuCUS) protocol. *Acad Emerg Med* 2015;**22**:182–91.
78. Čelutkienė J, Lainscak M, Anderson L, Gayat E, Grapsa J, Harjola VP et al. Imaging in patients with suspected acute heart failure: timeline approach position statement on behalf of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2020;**22**:181–95.
79. Hubert A, Girerd N, Le Breton H, Galli E, Latar I, Fournet M et al. Diagnostic accuracy of lung ultrasound for identification of elevated left ventricular filling pressure. *Int J Cardiol* 2019;**281**:62–8.
80. Miglioranza MH, Gargani L, Sant'anna RT, Rover MM, Martins VM, Mantovani A et al. Lung ultrasound for the evaluation of pulmonary congestion in outpatients: a comparison with clinical assessment, natriuretic peptides, and echocardiography. *JACC Cardiovasc Imaging* 2013;**6**:1141–51.
81. Kobayashi M, Gargani L, Palazzuoli A, Ambrosio G, Bayés-Genis A, Lupon J et al. Association between right-sided cardiac function and ultrasound-based pulmonary congestion on acutely decompensated heart failure: findings from a pooled analysis of four cohort studies. *Clin Res Cardiol* 2021;**110**:1181–92.
82. Neskovic AN, Skinner H, Price S, Via G, De Hert S, Stankovic I et al. Focus cardiac ultrasound core curriculum and core syllabus of the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2018;**19**:475–81.
83. Bedetti G, Gargani L, Corbisiero A, Frassi F, Poggianti E, Mottola G. Evaluation of ultrasound lung comets by hand-held echocardiography. *Cardiovasc Ultrasound* 2006;**4**:34.
84. Noble VE, Lamhaut L, Capp R, Bosson N, Liteplo A, Marx JS et al. Evaluation of a thoracic ultrasound training module for the detection of pneumothorax and pulmonary edema by prehospital physician care providers. *BMC Med Educ* 2009;**9**:3.
85. Zisis G, Yang Y, Huynh Q, Whitmore K, Lay M, Wright L et al. Nurse-provided lung and inferior vena cava assessment in patients with heart failure. *J Am Coll Cardiol* 2022;**80**:513–23.
86. de Souza LAM, Paredes RG, Giraldo T, Franco MH, de Carvalho-Filho MA, Cecilio-Fernandes D et al. Implementation and assessment of lung ultrasound training curriculum for physiotherapists with a focus on image acquisition and calculation of an aeration score. *Ultrasound Med Biol* 2022;**48**:2119–27.
87. Palazzuoli A, Ruocco G, Franci B, Evangelista I, Lucani B, Nuti R et al. Ultrasound indices of congestion in patients with acute heart failure according to body mass index. *Clin Res Cardiol* 2020;**109**:1423–33.